

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ZEPJULA safely and effectively. See full prescribing information for ZEPJULA.

ZEPJULA (niraparib) capsules, for oral use  
Initial U.S. Approval: 2017

### RECENT MAJOR CHANGES

Indications and Usage (1.1)	4/2020
Dosage and Administration (2.2)	4/2020
Dosage and Administration (2.3, 2.4)	3/2021
Warnings and Precautions (5.1, 5.3)	4/2020
Warnings and Precautions (5.2, 5.4, 5.6)	3/2021

### INDICATIONS AND USAGE

ZEPJULA is a poly (ADP-ribose) polymerase (PARP) inhibitor indicated:

- for the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to first-line platinum-based chemotherapy. (1.1)
- for the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy. (1.2)
- for the treatment of adult patients with advanced ovarian, fallopian tube, or primary peritoneal cancer who have been treated with 3 or more prior chemotherapy regimens and whose cancer is associated with homologous recombination deficiency (HRD) positive status defined by either:
  - a deleterious or suspected deleterious *BRCA* mutation, or
  - genomic instability and who have progressed more than 6 months after response to the last platinum-based chemotherapy.

Select patients for therapy based on an FDA-approved companion diagnostic for ZEPJULA. (1.3, 2.1)

### DOSAGE AND ADMINISTRATION

- First-line maintenance treatment of advanced ovarian cancer:**
  - For patients weighing <77 kg (<170 lbs) OR with a platelet count <150,000/mcL, the recommended dosage is 200 mg taken orally once daily. (2.2)
  - For patients weighing ≥77 kg (≥170 lbs) AND a platelet count ≥150,000/ mcL, the recommended dosage is 300 mg taken orally once daily. (2.2)
- For other indications:** The recommended dosage is 300 mg taken orally once daily. (2.2)
- Continue treatment until disease progression or unacceptable adverse reaction. (2.2)
- ZEPJULA may be taken with or without food. (2.2)
- For adverse reactions, consider interruption of treatment, dose reduction, or dose discontinuation. (2.3)

- For patients with moderate hepatic impairment, reduce the starting dosage of niraparib to 200 mg once daily. (2.4)

### DOSAGE FORMS AND STRENGTHS

Capsules: 100 mg (3)

### CONTRAINDICATIONS

None. (4)

### WARNINGS AND PRECAUTIONS

- Myelodysplastic Syndrome/Acute Myeloid Leukemia (MDS/AML):** MDS/AML occurred in patients exposed to ZEPJULA, and some cases were fatal. Monitor patients for hematological toxicity and discontinue if MDS/AML is confirmed. (5.1)
- Bone Marrow Suppression:** Test complete blood counts weekly for the first month, monthly for the next 11 months, and periodically thereafter for clinically significant changes. (5.2)
- Hypertension and Cardiovascular Effects:** Monitor blood pressure and heart rate at least weekly for the first 2 months, then monthly for the first year and periodically thereafter during treatment with ZEPJULA. Manage with antihypertensive medications and adjustment of the dose of ZEPJULA, if necessary. (5.3)
- Posterior Reversible Encephalopathy Syndrome (PRES):** PRES has occurred in patients treated with ZEPJULA. Discontinue ZEPJULA if PRES is confirmed. (5.4)
- Embryo-Fetal Toxicity:** ZEPJULA can cause fetal harm. Advise females of reproductive potential of the potential risk to a fetus and to use effective contraception. (5.5, 8.1, 8.3)
- Allergic Reactions to FD&C Yellow No. 5 (Tartrazine):** Contains FD&C Yellow No. 5 (tartrazine) as a color additive, which may cause allergic-type reactions (including bronchial asthma) in certain susceptible patients. (5.6)

### ADVERSE REACTIONS

Most common adverse reactions (incidence ≥10%) in patients who received ZEPJULA were nausea, thrombocytopenia, anemia, fatigue, constipation, musculoskeletal pain, abdominal pain, vomiting, neutropenia, decreased appetite, leukopenia, insomnia, headache, dyspnea, rash, diarrhea, hypertension, cough, dizziness, acute kidney injury, urinary tract infection, and hypomagnesemia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact GlaxoSmithKline at 1-888-825-5249 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### USE IN SPECIFIC POPULATIONS

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 3/2021

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

- 1.1 First-Line Maintenance Treatment of Advanced Ovarian Cancer
- 1.2 Maintenance Treatment of Recurrent Ovarian Cancer
- 1.3 Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Patient Selection for Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies
- 2.2 Recommended Dosage
- 2.3 Dosage Adjustments for Adverse Reactions
- 2.4 Dosage Adjustment for Hepatic Impairment

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Myelodysplastic Syndrome/Acute Myeloid Leukemia
- 5.2 Bone Marrow Suppression
- 5.3 Hypertension and Cardiovascular Effects
- 5.4 Posterior Reversible Encephalopathy Syndrome
- 5.5 Embryo-Fetal Toxicity
- 5.6 Allergic Reactions to FD&C Yellow No. 5 (Tartrazine)

### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

6.2 Postmarketing Experience

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

### 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

### 14 CLINICAL STUDIES

- 14.1 First-Line Maintenance Treatment of Advanced Ovarian Cancer
- 14.2 Maintenance Treatment of Recurrent Ovarian Cancer
- 14.3 Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies

---

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

#### 1.1 First-Line Maintenance Treatment of Advanced Ovarian Cancer

ZEJULA is indicated for the maintenance treatment of adult patients with advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to first-line platinum-based chemotherapy.

#### 1.2 Maintenance Treatment of Recurrent Ovarian Cancer

ZEJULA is indicated for the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy.

#### 1.3 Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies

ZEJULA is indicated for the treatment of adult patients with advanced ovarian, fallopian tube, or primary peritoneal cancer who have been treated with 3 or more prior chemotherapy regimens and whose cancer is associated with homologous recombination deficiency (HRD) positive status defined by either:

- a deleterious or suspected deleterious *BRCA* mutation, or
- genomic instability and who have progressed more than 6 months after response to the last platinum-based chemotherapy [*see Clinical Studies (14.3)*].

Select patients for therapy based on an FDA-approved companion diagnostic for ZEJULA.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Patient Selection for Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies

Select patients for treatment of advanced ovarian cancer after 3 or more chemotherapy regimens associated with HRD positive status based on either deleterious or suspected deleterious *BRCA* mutation and/or genomic instability score (GIS) [*see Clinical Studies (14.3)*].

Information on FDA-approved tests for the detection of either deleterious or suspected deleterious *BRCA* mutation or genomic instability for this indication is available at <https://www.fda.gov/companiondiagnostics>.

#### 2.2 Recommended Dosage

Continue treatment with ZEJULA until disease progression or unacceptable toxicity.

Instruct patients to take their dose of ZEJULA at approximately the same time each day. Advise patients to swallow each capsule whole and not to chew, crush, or split ZEJULA prior to

swallowing. ZEPJULA may be taken with or without food. Bedtime administration may be a potential method for managing nausea.

In the case of a missed dose of ZEPJULA, instruct patients to take their next dose at its regularly scheduled time. If a patient vomits or misses a dose of ZEPJULA, an additional dose should not be taken.

#### First-Line Maintenance Treatment of Advanced Ovarian Cancer

- For patients weighing <77 kg (<170 lbs) OR with a platelet count of <150,000/mcL, the recommended dosage is 200 mg (two 100-mg capsules) taken orally once daily.
- For patients weighing ≥77 kg (≥170 lbs) AND who have a platelet count ≥150,000/mcL, the recommended dosage is 300 mg (three 100-mg capsules) taken orally once daily.

For the maintenance treatment of advanced ovarian cancer, patients should start treatment with ZEPJULA no later than 12 weeks after their most recent platinum-containing regimen.

#### Maintenance Treatment of Recurrent Ovarian Cancer

The recommended dosage of ZEPJULA is 300 mg (three 100-mg capsules) taken orally once daily.

For the maintenance treatment of recurrent ovarian cancer, patients should start treatment with ZEPJULA no later than 8 weeks after their most recent platinum-containing regimen.

#### Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies

The recommended dosage of ZEPJULA is 300 mg (three 100-mg capsules) taken orally once daily.

### **2.3 Dosage Adjustments for Adverse Reactions**

To manage adverse reactions, consider interruption of treatment, dose reduction, or dose discontinuation. The recommended dose modifications for adverse reactions are listed in Tables 1, 2, and 3.

**Table 1. Recommended Dose Modifications for Adverse Reactions**

<b>Starting Dose Level</b>	<b>200 mg</b>	<b>300 mg</b>
First dose reduction	100 mg/day <sup>a</sup> (one 100-mg capsule)	200 mg/day (two 100-mg capsules)
Second dose reduction	Discontinue ZEPJULA.	100 mg/day <sup>a</sup> (one 100-mg capsule)

<sup>a</sup> If further dose reduction below 100 mg/day is required, discontinue ZEPJULA.

**Table 2. Dose Modifications for Non-Hematologic Adverse Reactions**

Non-hematologic CTCAE $\geq$ Grade 3 adverse reaction that persists despite medical management	<ul style="list-style-type: none"> <li>Withhold ZEPJULA for a maximum of 28 days or until resolution of adverse reaction.</li> <li>Resume ZEPJULA at a reduced dose per Table 1.</li> </ul>
CTCAE $\geq$ Grade 3 treatment-related adverse reaction lasting more than 28 days while patient is administered ZEPJULA 100 mg/day	Discontinue ZEPJULA.

CTCAE = Common Terminology Criteria for Adverse Events.

**Table 3. Dose Modifications for Hematologic Adverse Reactions**

Monitor complete blood counts weekly for the first month, monthly for the next 11 months of treatment, and periodically after this time [see <i>Warnings and Precautions (5.1)</i> ].	
Platelet count $<100,000/\text{mL}$	<p>First occurrence:</p> <ul style="list-style-type: none"> <li>Withhold ZEPJULA for a maximum of 28 days and monitor blood counts weekly until platelet counts return to <math>\geq 100,000/\text{mL}</math>.</li> <li>Resume ZEPJULA at same or reduced dose per Table 1.</li> <li>If platelet count is <math>&lt;75,000/\text{mL}</math>, resume at a reduced dose.</li> </ul> <p>Second occurrence:</p> <ul style="list-style-type: none"> <li>Withhold ZEPJULA for a maximum of 28 days and monitor blood counts weekly until platelet counts return to <math>\geq 100,000/\text{mL}</math>.</li> <li>Resume ZEPJULA at a reduced dose per Table 1.</li> <li>Discontinue ZEPJULA if the platelet count has not returned to acceptable levels within 28 days of the dose interruption period or if the patient has already undergone dose reduction to 100 mg once daily.<sup>a</sup></li> </ul>
Neutrophil $<1,000/\text{mL}$ or hemoglobin $<8 \text{ g/dL}$	<ul style="list-style-type: none"> <li>Withhold ZEPJULA for a maximum of 28 days and monitor blood counts weekly until neutrophil counts return to <math>\geq 1,500/\text{mL}</math> or hemoglobin returns to <math>\geq 9 \text{ g/dL}</math>.</li> <li>Resume ZEPJULA at a reduced dose per Table 1.</li> <li>Discontinue ZEPJULA if neutrophils and/or hemoglobin have not returned to acceptable levels within 28 days of the dose interruption period or if the patient has already undergone dose reduction to 100 mg once daily.<sup>a</sup></li> </ul>

Hematologic adverse reaction requiring transfusion	<ul style="list-style-type: none"> <li>• For patients with platelet count <math>\leq 10,000/\text{mcL}</math>, platelet transfusion should be considered. If there are other risk factors such as coadministration of anticoagulation or antiplatelet drugs, consider interrupting these drugs and/or transfusion at a higher platelet count.</li> <li>• Resume ZEJULA at a reduced dose.</li> </ul>
--	--

<sup>a</sup> If myelodysplastic syndrome or acute myeloid leukemia (MDS/AML) is confirmed, discontinue ZEJULA [see *Warnings and Precautions (5.1, 5.2)*].

## 2.4 Dosage Adjustment for Hepatic Impairment

### Moderate Hepatic Impairment

For patients with moderate hepatic impairment, reduce the starting dosage of ZEJULA to 200 mg once daily. Monitor patients for hematologic toxicity and reduce the dose further, if needed [see *Dosage and Administration (2.3)*, *Use in Specific Populations (8.7)*, *Clinical Pharmacology (12.3)*].

## 3 DOSAGE FORMS AND STRENGTHS

100-mg capsule having a white body with “100 mg” printed in black ink, and a purple cap with “Niraparib” printed in white ink.

## 4 CONTRAINDICATIONS

None.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Myelodysplastic Syndrome/Acute Myeloid Leukemia

Myelodysplastic syndrome/acute myeloid leukemia (MDS/AML), including cases with fatal outcome, have been reported in patients who received monotherapy with ZEJULA in clinical trials. In 1,785 patients treated with ZEJULA in clinical trials, MDS/AML occurred in 15 patients (0.8%).

The duration of therapy with ZEJULA in patients who developed secondary MDS/cancer therapy-related AML varied from 0.5 months to 4.9 years. All of these patients had received previous chemotherapy with platinum agents and/or other DNA-damaging agents, including radiotherapy. Discontinue ZEJULA if MDS/AML is confirmed.

### 5.2 Bone Marrow Suppression

Hematologic adverse reactions, including thrombocytopenia, anemia, neutropenia, and/or pancytopenia have been reported in patients treated with ZEJULA [see *Adverse Reactions (6)*].

In PRIMA, the overall incidences of  $\geq$ Grade 3 thrombocytopenia, anemia, and neutropenia were reported in 39%, 31%, and 21%, respectively, of patients receiving ZEJULA. Discontinuation due to thrombocytopenia, anemia, and neutropenia occurred in 4%, 2%, and 2%, respectively, of patients. In patients who were administered a starting dose of ZEJULA based on baseline weight or platelet count,  $\geq$ Grade 3 thrombocytopenia, anemia, and neutropenia were reported in 22%, 23%, and 15%, respectively, of patients receiving ZEJULA. Discontinuation due to thrombocytopenia, anemia, and neutropenia occurred in 3%, 3%, and 2%, respectively, of patients.

In NOVA,  $\geq$ Grade 3 thrombocytopenia, anemia, and neutropenia were reported in 29%, 25%, and 20%, respectively, of patients receiving ZEJULA. Discontinuation due to thrombocytopenia, anemia, and neutropenia occurred in 3%, 1%, and 2%, respectively, of patients.

In QUADRA,  $\geq$ Grade 3 thrombocytopenia, anemia, and neutropenia were reported in 28%, 27%, and 13%, respectively, of patients receiving ZEJULA. Discontinuation due to thrombocytopenia, anemia, and neutropenia occurred in 4%, 2%, and 1%, respectively, of patients.

Do not start ZEJULA until patients have recovered from hematological toxicity caused by previous chemotherapy ( $\leq$ Grade 1). Monitor complete blood counts weekly for the first month, monthly for the next 11 months of treatment, and periodically after this time. If hematological toxicities do not resolve within 28 days following interruption, discontinue ZEJULA and refer the patient to a hematologist for further investigations, including bone marrow analysis and blood sample for cytogenetics [*see Dosage and Administration (2.3)*].

### **5.3 Hypertension and Cardiovascular Effects**

Hypertension and hypertensive crisis have been reported in patients treated with ZEJULA.

In PRIMA, Grade 3 to 4 hypertension occurred in 6% of patients treated with ZEJULA compared with 1% of placebo-treated patients with a median time from first dose to first onset of 43 days (range: 1 to 531 days) and with a median duration of 12 days (range: 1 to 61 days). There were no discontinuations due to hypertension.

In NOVA, Grade 3 to 4 hypertension occurred in 9% of patients treated with ZEJULA compared with 2% of placebo-treated patients with a median time from first dose to first onset of 77 days (range: 4 to 504 days) and with a median duration of 15 days (range: 1 to 86 days). Discontinuation due to hypertension occurred in  $<1\%$  of patients.

In QUADRA, Grade 3 to 4 hypertension occurred in 5% of patients treated with ZEJULA with a median time from first dose to first onset of 15 days (range: 1 to 316 days) and with a median duration of 7 days (range: 1 to 118 days). Discontinuation due to hypertension occurred in  $<0.2\%$  of patients.

Monitor blood pressure and heart rate at least weekly for the first 2 months, then monthly for the first year and periodically thereafter during treatment with ZEJULA. Closely monitor patients with cardiovascular disorders, especially coronary insufficiency, cardiac arrhythmias, and

hypertension. Medically manage hypertension with antihypertensive medications and adjustment of the dose of ZEJULA, if necessary [see *Dosage and Administration (2.3)*, *Nonclinical Toxicology (13.2)*].

#### **5.4 Posterior Reversible Encephalopathy Syndrome**

Posterior reversible encephalopathy syndrome (PRES) occurred in 0.1% of 2,165 patients treated with ZEJULA in clinical trials and has also been described in postmarketing reports [see *Adverse Reactions (6.2)*]. Signs and symptoms of PRES include seizure, headache, altered mental status, visual disturbance, or cortical blindness, with or without associated hypertension. A diagnosis of PRES requires confirmation by brain imaging, preferably magnetic resonance imaging.

Monitor all patients treated with ZEJULA for signs and symptoms of PRES. If PRES is suspected, promptly discontinue ZEJULA and administer appropriate treatment. The safety of reinitiating ZEJULA in patients previously experiencing PRES is not known.

#### **5.5 Embryo-Fetal Toxicity**

Based on its mechanism of action, ZEJULA can cause fetal harm when administered to a pregnant woman [see *Clinical Pharmacology (12.1)*]. ZEJULA has the potential to cause teratogenicity and/or embryo-fetal death since niraparib is genotoxic and targets actively dividing cells in animals and patients (e.g., bone marrow) [see *Warnings and Precautions (5.2)*, *Nonclinical Toxicology (13.1)*]. Due to the potential risk to a fetus based on its mechanism of action, animal developmental and reproductive toxicology studies were not conducted with niraparib.

Apprise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment and for 6 months after the last dose of ZEJULA [see *Use in Specific Populations (8.1, 8.3)*].

#### **5.6 Allergic Reactions to FD&C Yellow No. 5 (Tartrazine)**

ZEJULA capsules contain FD&C Yellow No. 5 (tartrazine), which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons. Although the overall incidence of FD&C Yellow No. 5 (tartrazine) sensitivity in the general population is low, it is frequently seen in patients who also have aspirin hypersensitivity.

### **6 ADVERSE REACTIONS**

The following clinically significant adverse reactions are described elsewhere in the labeling:

- MDS/AML [see *Warnings and Precautions (5.1)*]
- Bone marrow suppression [see *Warnings and Precautions (5.2)*]
- Hypertension and cardiovascular effects [see *Warnings and Precautions (5.3)*]
- Posterior reversible encephalopathy syndrome [see *Warnings and Precautions (5.4)*]

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The most common adverse reactions of all grades in >10% of 1,314 patients who received ZEPJULA in the pooled PRIMA, NOVA, and QUADRA trials were nausea (65%), thrombocytopenia (60%), anemia (56%), fatigue (55%), constipation (39%), musculoskeletal pain (36%), abdominal pain (35%), vomiting (33%), neutropenia (31%), decreased appetite (24%), leukopenia (24%), insomnia (23%), headache (23%), dyspnea (22%), rash (21%), diarrhea (18%), hypertension (17%), cough (16%), dizziness (14%), acute kidney injury (13%), urinary tract infection (12%), and hypomagnesemia (11%).

### First-Line Maintenance Treatment of Advanced Ovarian Cancer

The safety of ZEPJULA for the treatment of patients with advanced ovarian cancer following first-line treatment with platinum-based chemotherapy was studied in the PRIMA trial, a placebo-controlled, double-blind study in which 728 patients received niraparib or placebo. Among patients who received ZEPJULA, the median duration of treatment was 11.1 months (range: 0.03 to 29 months).

*All Patients Receiving ZEPJULA in PRIMA:* Serious adverse reactions occurred in 32% of patients receiving ZEPJULA. Serious adverse reactions in >2% of patients were thrombocytopenia (16%), anemia (6%), and small intestinal obstruction (2.9%). Fatal adverse reactions occurred in 0.4% of patients, including intestinal perforation and pleural effusion (1 patient each).

Permanent discontinuation due to adverse reactions occurred in 12% of patients who received ZEPJULA. Adverse reactions resulting in permanent discontinuation in >1% of patients who received ZEPJULA included thrombocytopenia (3.7%), anemia (1.9%), and nausea and neutropenia (1.2% each).

Adverse reactions led to dose reduction or interruption in 80% of patients, most frequently from thrombocytopenia (56%), anemia (33%), and neutropenia (20%).

Table 4 and Table 5 summarize the common adverse reactions and abnormal laboratory findings, respectively, observed in all patients treated with ZEPJULA in the PRIMA study.



**Table 4. Adverse Reactions Reported in  $\geq 10\%$  of All Patients Receiving ZEJULA in PRIMA<sup>a</sup>**

Adverse Reaction	Grades 1-4 <sup>b</sup>		Grades 3-4 <sup>b</sup>	
	ZEJULA (n = 484) %	Placebo (n = 244) %	ZEJULA (n = 484) %	Placebo (n = 244) %
Blood and lymphatic system disorders				
Thrombocytopenia	66	5	39	0.4
Anemia	64	18	31	2
Neutropenia <sup>c</sup>	42	8	21	1
Leukopenia <sup>d</sup>	28	9	5	0.4
Gastrointestinal disorders				
Nausea	57	28	1	1
Constipation	40	20	1	0.4
Vomiting	22	12	1	1
General disorders and administration site conditions				
Fatigue	51	41	3	1
Investigations				
AST/ALT elevation	14	7	3	0.8
Metabolism and nutrition disorders				
Decreased appetite	19	8	1	0
Musculoskeletal and connective tissue disorders				
Musculoskeletal pain	39	38	1	0
Nervous system disorders				
Headache	26	15	0.4	0
Dizziness	19	13	0	0.4
Psychiatric disorders				
Insomnia	25	15	1	0.4
Renal and urinary disorders				
Acute kidney injury <sup>e</sup>	12	5	0.2	0
Respiratory, thoracic, and mediastinal disorders				
Dyspnea	22	13	0.4	1
Cough	18	15	0	0.4
Vascular disorders				
Hypertension	18	7	6	1

AST/ALT = Aspartate transaminase/alanine aminotransferase.

<sup>a</sup> All adverse reactions in the table consist of grouped preferred terms except for nausea, vomiting, decreased appetite, headache, and insomnia, which are single preferred terms.

<sup>b</sup> Common Terminology Criteria for Adverse Events version 4.02.

<sup>c</sup> Includes neutropenia, neutropenic infection, neutropenic sepsis, and febrile neutropenia.

<sup>d</sup> Includes leukopenia, lymphocyte count decreased, lymphopenia, and white blood cell count decreased.

<sup>e</sup> Includes blood creatinine increased, blood urea increased, acute kidney injury, renal failure, and blood creatine increased.

**Table 5. Abnormal Laboratory Findings in  $\geq 25\%$  of All Patients Receiving ZEJULA in PRIMA**

Abnormal Laboratory Finding	Grades 1-4		Grades 3-4	
	ZEJULA (n = 484) %	Placebo (n = 244) %	ZEJULA (n = 484) %	Placebo (n = 244) %
Decreased hemoglobin	87	66	29	1
Decreased platelets	74	13	37	0
Decreased leukocytes	71	36	9	0
Increased glucose	66	57	3	3
Decreased neutrophils	66	25	23	1
Decreased lymphocytes	51	29	7	3
Increased alkaline phosphatase	46	21	1	0
Increased creatinine	40	23	0	0
Decreased magnesium	36	34	1	0
Increased aspartate aminotransferase	35	17	1	0.4
Increased alanine aminotransferase	29	17	2	1

*Patients Receiving ZEJULA with Dose Based on Baseline Weight or Platelet Count in PRIMA:* Among patients who received ZEJULA with the dose based on weight and platelet count, the median duration of treatment was 11 months (range: 1 day to 16 months).

Serious adverse reactions occurred in 27% of patients receiving ZEJULA. Serious adverse reactions in  $>2\%$  of patients were anemia (8%), and thrombocytopenia (7%). No fatal adverse reactions occurred.

Permanent discontinuation due to adverse reactions occurred in 14% of patients who received ZEJULA. Adverse reactions resulting in permanent discontinuation in  $>2\%$  of patients who received ZEJULA included thrombocytopenia and anemia (3% each) and nausea (2.4%).

Adverse reactions led to dose reduction or interruption in 72% of patients, most frequently from thrombocytopenia (40%), anemia (23%), and neutropenia (15%).

Table 6 and Table 7 summarize adverse reactions and abnormal laboratory findings in the group

of patients who received ZEJULA.

**Table 6. Adverse Reactions Reported in  $\geq 10\%$  of Patients Receiving ZEJULA Based on Baseline Weight or Platelet Count in PRIMA<sup>a</sup>**

Adverse Reaction	Grades 1-4 <sup>b</sup>		Grades 3-4 <sup>b</sup>	
	ZEJULA (n = 169) %	Placebo (n = 86) %	ZEJULA (n = 169) %	Placebo (n = 86) %
Blood and lymphatic system disorders				
Thrombocytopenia	54	5	21	1
Anemia	50	28	23	1
Neutropenia <sup>c</sup>	36	8	15	1
Leukopenia <sup>d</sup>	28	11	5	0
Gastrointestinal disorders				
Nausea	53	21	1	0
Constipation	31	15	1	1
Vomiting	17	9	0	1
General disorders and administration site conditions				
Fatigue	48	36	3	0
Metabolism and nutrition disorders				
Decreased appetite	19	5	1	0
Nervous system disorders				
Headache	22	17	1	0
Dizziness	14	13	0	0
Psychiatric disorders				
Insomnia	21	14	0	0
Renal and urinary disorders				
Acute kidney injury <sup>e</sup>	21	14	0	0
Respiratory, thoracic, and mediastinal disorders				
Dyspnea	18	10	0	1
Vascular disorders				
Hypertension	17	9	5	2

<sup>a</sup> All adverse reactions in the table consist of grouped preferred terms except for nausea, vomiting, decreased appetite, headache, and insomnia, which are single preferred terms.

<sup>b</sup> Common Terminology Criteria for Adverse Events version 4.02.

<sup>c</sup> Includes neutropenia, neutropenic infection, neutropenic sepsis, and febrile neutropenia.

<sup>d</sup> Includes leukopenia, lymphocyte count decreased, lymphopenia, and white blood cell count decreased.

<sup>e</sup> Includes blood creatinine increased, blood urea increased, acute kidney injury, renal failure, and blood creatine increased.

**Table 7. Abnormal Laboratory Findings in  $\geq 25\%$  of All Patients Receiving ZEJULA Based on Baseline Weight or Platelet Count in PRIMA**

Abnormal Laboratory Finding	Grades 1-4		Grades 3-4	
	ZEJULA (n = 169) %	Placebo (n = 86) %	ZEJULA (n = 169) %	Placebo (n = 86) %
Decreased hemoglobin	81	70	21	0
Decreased leukocytes	70	36	6	0
Decreased platelets	63	15	18	0
Increased glucose	63	56	2	1
Decreased neutrophils	60	27	15	0
Decreased lymphocytes	52	30	5	4
Increased alkaline phosphatase	43	17	1	0
Decreased magnesium	44	30	0	0
Increased creatinine	41	22	0	0
Increased aspartate aminotransferase	31	19	1	0
Increased alanine aminotransferase	28	15	2	2

#### Maintenance Treatment of Recurrent Ovarian Cancer

The safety of monotherapy with ZEJULA 300 mg once daily has been studied in 367 patients with platinum-sensitive recurrent ovarian, fallopian tube, and primary peritoneal cancer in the NOVA trial. Adverse reactions in NOVA led to dose reduction or interruption in 69% of patients, most frequently from thrombocytopenia (41%) and anemia (20%). The permanent discontinuation rate due to adverse reactions in NOVA was 15%. The median exposure to ZEJULA in these patients was 250 days.

Table 8 and Table 9 summarize the common adverse reactions and abnormal laboratory findings, respectively, observed in patients treated with ZEJULA in NOVA.

**Table 8. Adverse Reactions Reported in ≥10% of Patients Receiving ZEPJULA in NOVA**

Adverse Reaction	Grades 1-4 <sup>a</sup>		Grades 3-4 <sup>a</sup>	
	ZEPJULA (n = 367) %	Placebo (n = 179) %	ZEPJULA (n = 367) %	Placebo (n = 179) %
Blood and lymphatic system disorders				
Thrombocytopenia	61	5	29	0.6
Anemia	50	7	25	0
Neutropenia <sup>b</sup>	30	6	20	2
Leukopenia	17	8	5	0
Cardiac disorders				
Palpitations	10	2	0	0
Gastrointestinal disorders				
Nausea	74	35	3	1
Constipation	40	20	0.8	2
Vomiting	34	16	2	0.6
Mucositis/stomatitis	20	6	0.5	0
Dyspepsia	18	12	0	0
Dry mouth	10	4	0.3	0
General disorders and administration site conditions				
Fatigue/asthenia	57	41	8	0.6
Metabolism and nutrition disorders				
Decreased appetite	25	15	0.3	0.6
Infections and infestations				
Urinary tract infection	13	8	0.8	1
Investigations				
AST/ALT elevation	10	5	4	2
Musculoskeletal and connective tissue disorders				
Back pain	18	12	0.8	0
Nervous system disorders				
Headache	26	11	0.3	0
Dizziness	18	8	0	0
Dysgeusia	10	4	0	0
Psychiatric disorders				
Insomnia	27	8	0.3	0
Anxiety	11	7	0.3	0.6

Respiratory, thoracic, and mediastinal disorders				
Nasopharyngitis	23	14	0	0
Dyspnea	20	8	1	1
Cough	16	5	0	0
Skin and subcutaneous tissue disorders				
Rash	21	9	0.5	0
Vascular disorders				
Hypertension	20	5	9	2

AST/ALT = Aspartate transaminase/alanine aminotransferase.

<sup>a</sup> Common Terminology Criteria for Adverse Events version 4.02.

<sup>b</sup> Includes preferred terms of neutropenic infection, neutropenic sepsis, and febrile neutropenia.

**Table 9. Abnormal Laboratory Findings in  $\geq 25\%$  of Patients Receiving ZEJULA in NOVA**

Abnormal Laboratory Finding	Grades 1-4		Grades 3-4	
	ZEJULA (n = 367)	Placebo (n = 179)	ZEJULA (n = 367)	Placebo (n = 179)
	%	%	%	%
Decrease in hemoglobin	85	56	25	0.5
Decrease in platelet count	72	21	35	0.5
Decrease in white blood cell count	66	37	7	0.7
Decrease in absolute neutrophil count	53	25	21	2
Increase in aspartate aminotransferase	36	23	1	0
Increase in alanine aminotransferase	28	15	1	2

The following adverse reactions and laboratory abnormalities have been identified in  $\geq 1$  to  $< 10\%$  of the 367 patients receiving ZEJULA in the NOVA trial and not included in the table:

tachycardia, peripheral edema, hypokalemia, bronchitis, conjunctivitis, gamma-glutamyl transferase increased, blood creatinine increased, blood alkaline phosphatase increased, weight decreased, depression, and epistaxis.

#### Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies

The safety of monotherapy with ZEJULA 300 mg once daily has been studied in QUADRA, a single-arm study in 463 patients with recurrent high-grade serous epithelial ovarian, fallopian tube, or primary peritoneal cancer who had been treated with 3 or more prior lines of therapy. The median duration of overall study treatment was 3 months (range: 0.03 to 32 months). For the indicated QUADRA population, the median duration was 4 months (range: 0.1 to 30 months).

Fatal adverse reactions occurred in 2% of patients, including cardiac arrest.

Serious adverse reactions occurred in 43% of patients receiving ZEJULA. Serious adverse reactions in >3% of patients were small intestinal obstruction (7%), vomiting (6%), nausea (5%), and abdominal pain (4%).

Permanent discontinuation due to adverse reactions (Grade 1 to 4) occurred in 21% of patients who received ZEJULA.

Adverse reactions led to dose reduction or interruption in 73% of patients receiving ZEJULA. The most common adverse reactions ( $\geq 5\%$ ) resulting in dose reduction or interruption of ZEJULA were thrombocytopenia (40%), anemia (21%), neutropenia (11%), nausea (13%), vomiting (11%), fatigue (9%), and abdominal pain (5%).

Table 10 and Table 11 summarize the common adverse reactions and abnormal laboratory findings, respectively, observed in patients treated with ZEJULA in QUADRA.

**Table 10. Adverse Reactions Reported in  $\geq 10\%$  of Patients Receiving ZEJULA in QUADRA**

<b>Adverse Reaction</b>	<b>Grades 1-4<sup>a</sup> (n = 463) %</b>	<b>Grades 3-4<sup>a</sup> (n = 463) %</b>
Blood and lymphatic system disorders		
Anemia <sup>b</sup>	51	27
Thrombocytopenia <sup>c</sup>	52	28
Neutropenia <sup>d</sup>	20	13
Gastrointestinal disorders		
Nausea	67	10
Vomiting	44	8
Constipation	36	5
Abdominal pain	34	7
Diarrhea	17	0.2
General disorders and administration site conditions		
Fatigue	56	7
Infections and infestations		
Urinary tract infection	15	2
Investigations		
Blood alkaline phosphatase increased	11	2
AST/ALT elevation	11	1
Metabolism and nutrition disorders		
Decreased appetite	27	2

Musculoskeletal and connective tissue disorders		
Musculoskeletal pain	29	3
Nervous system disorders		
Headache	19	0.4
Dizziness	11	0
Psychiatric disorders		
Insomnia	21	1
Renal and urinary disorders		
Acute kidney injury	17	1
Respiratory, thoracic, and mediastinal disorders		
Dyspnea	22	3
Cough	13	0
Vascular disorders		
Hypertension	14	5

AST/ALT = Aspartate transaminase/alanine aminotransferase.

<sup>a</sup> Common Terminology Criteria for Adverse Events version 4.02.

<sup>b</sup> Anemia includes events with preferred terms of anemia, hemoglobin decreased, anemia macrocytic, aplastic anemia, and normochromic normocytic anemia.

<sup>c</sup> Thrombocytopenia includes events with preferred terms of thrombocytopenia and platelet count decreased.

<sup>d</sup> Neutropenia includes events with preferred terms of neutropenia, neutrophil count decreased, neutropenic infection, and neutropenic sepsis.

**Table 11. Abnormal Laboratory Findings in  $\geq 25\%$  of Patients Receiving ZEJULA in QUADRA**

Abnormal Laboratory Finding	Grades 1-4 (n = 463)	Grades 3-4 (n = 463)
	%	%
Decreased hemoglobin	83	26
Increased glucose	66	5
Decreased platelets	60	28
Decreased lymphocytes	57	18
Decreased leukocytes	53	9
Decreased magnesium	46	1
Increased alkaline phosphatase	40	4
Increased gamma glutamyl transferase	40	8
Increased creatinine	36	0.4
Decreased sodium	34	6



Decreased neutrophils	34	15
Increased aspartate aminotransferase	29	2
Decreased albumin	27	2

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of ZEPJULA. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

### Blood and Lymphatic System Disorders

Pancytopenia.

### Immune System Disorders

Hypersensitivity (including anaphylaxis).

### Nervous System Disorders

Posterior reversible encephalopathy syndrome (PRES).

### Psychiatric Disorders

Confusional state/disorientation, hallucination, cognitive impairment (e.g., memory impairment, concentration impairment).

### Respiratory, Thoracic, and Mediastinal Disorders

Non-infectious pneumonitis.

### Skin and Subcutaneous Tissue Disorders

Photosensitivity.

### Vascular Disorders

Hypertensive crisis.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Based on its mechanism of action, ZEPJULA can cause fetal harm when administered to pregnant women [see *Clinical Pharmacology (12.1)*]. There are no data regarding the use of ZEPJULA in pregnant women to inform the drug-associated risk. ZEPJULA has the potential to cause teratogenicity and/or embryo-fetal death since niraparib is genotoxic and targets actively dividing cells in animals and patients (e.g., bone marrow) [see *Warnings and Precautions (5.2)*, *Nonclinical Toxicology (13.1)*]. Due to the potential risk to a fetus based on its mechanism of

action, animal developmental and reproductive toxicology studies were not conducted with niraparib. Advise pregnant women of the potential risk to a fetus.

The background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## **8.2 Lactation**

### Risk Summary

No data are available regarding the presence of niraparib or its metabolites in human milk, or on its effects on the breastfed child or milk production. Because of the potential for serious adverse reactions in a breastfed child, advise a lactating woman not to breastfeed during treatment with ZEJULA and for 1 month after receiving the final dose.

## **8.3 Females and Males of Reproductive Potential**

ZEJULA can cause fetal harm when administered to a pregnant woman [*see Use in Specific Populations (8.1)*].

### Pregnancy Testing

Verify the pregnancy status of females of reproductive potential prior to initiating treatment with ZEJULA.

### Contraception

*Females:* Advise females of reproductive potential to use effective contraception during treatment with ZEJULA and for at least for 6 months following the last dose.

### Infertility

*Males:* Based on animal studies, ZEJULA may impair fertility in males of reproductive potential [*see Nonclinical Toxicology (13.1)*].

## **8.4 Pediatric Use**

The safety and effectiveness of ZEJULA have not been established in pediatric patients.

## **8.5 Geriatric Use**

In PRIMA, 39% of patients were aged 65 years or older and 10% were aged 75 years or older. In NOVA, 35% of patients were aged 65 years or older and 8% were aged 75 years or older. No overall differences in safety and effectiveness of ZEJULA were observed between these patients and younger patients but greater sensitivity of some older individuals cannot be ruled out.

## **8.6 Renal Impairment**

No dose adjustment is necessary for patients with mild (CLCr: 60 to 89 mL/min) to moderate (CLCr: 30 to 59 mL/min) renal impairment. The degree of renal impairment was determined by

creatinine clearance as estimated by the Cockcroft-Gault equation. The safety of ZEJULA in patients with severe renal impairment or end-stage renal disease undergoing hemodialysis is unknown.

## 8.7 Hepatic Impairment

For patients with moderate hepatic impairment, reduce the starting dosage of niraparib to 200 mg once daily [see *Dosage and Administration* (2.4)]. Niraparib exposure increased in patients with moderate hepatic impairment [total bilirubin  $\geq 1.5$  x upper level of normal (ULN) to 3.0 x ULN and any aspartate transaminase (AST) level]. Monitor patients for hematologic toxicity and reduce the dose further, if needed [see *Dosage and Administration* (2.3)].


For patients with mild hepatic impairment (total bilirubin  $< 1.5$  x ULN and any AST level or bilirubin  $\leq$  ULN and AST  $>$  ULN), no dose adjustment is needed.

The recommended dose of ZEJULA has not been established for patients with severe hepatic impairment (total bilirubin  $> 3.0$  x ULN and any AST level) [see *Clinical Pharmacology* (12.3)].

## 11 DESCRIPTION

Niraparib is an orally available poly (ADP-ribose) polymerase (PARP) inhibitor.

The chemical name for niraparib tosylate monohydrate is 2-{4-[(3S)-piperidin-3-yl]phenyl}-2H-indazole 7-carboxamide 4-methylbenzenesulfonate hydrate (1:1:1). The molecular formula is  $C_{26}H_{30}N_4O_5S$  and it has a molecular weight of 510.61 amu. The molecular structure is shown below:



Niraparib tosylate monohydrate is a white to off-white, non-hygroscopic crystalline solid.

Niraparib solubility is pH independent below the pKa of 9.95, with an aqueous free base solubility of 0.7 mg/mL to 1.1 mg/mL across the physiological pH range.

Each ZEJULA capsule contains 159.4 mg niraparib tosylate monohydrate equivalent to 100 mg niraparib free base as the active ingredient. The inactive ingredients in the *capsule fill* are magnesium stearate and lactose monohydrate. The *capsule shell* consists of titanium dioxide and gelatin in the white capsule body and FD&C Blue No. 1, FD&C Red No. 3, FD&C Yellow No. 5 (tartrazine), and gelatin in the purple capsule cap. The *black printing ink* consists of shellac, dehydrated alcohol, isopropyl alcohol, butyl alcohol, propylene glycol, purified water, strong ammonia solution, potassium hydroxide, and black iron oxide. The *white printing ink* consists of shellac, dehydrated alcohol, isopropyl alcohol, butyl alcohol, propylene glycol, sodium hydroxide, povidone, and titanium dioxide.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Niraparib is an inhibitor of PARP enzymes, including PARP-1 and PARP-2, that play a role in DNA repair. In vitro studies have shown that niraparib-induced cytotoxicity may involve inhibition of PARP enzymatic activity and increased formation of PARP-DNA complexes resulting in DNA damage, apoptosis, and cell death. Increased niraparib-induced cytotoxicity was observed in tumor cell lines with or without deficiencies in *BRCA1/2*. Niraparib decreased tumor growth in mouse xenograft models of human cancer cell lines with deficiencies in *BRCA1/2* and in human patient-derived xenograft tumor models with HRD that had either mutated or wild-type *BRCA1/2*.

### 12.2 Pharmacodynamics

The pharmacodynamic response of niraparib has not been characterized.

#### Hypertension and Cardiovascular Effects

Niraparib has the potential to cause effects on pulse rate and blood pressure in patients receiving the recommended dose, which may be related to pharmacological inhibition of the dopamine transporter (DAT), norepinephrine transporter (NET), and serotonin transporter (SERT) [*see Nonclinical Toxicology (13.2)*].

In the PRIMA study, mean pulse rate and blood pressure increased over baseline in the niraparib arm relative to the placebo arm at most on-study assessments. Mean greatest increases from baseline in pulse rate on treatment were 22.4 and 14.0 beats/min in the niraparib and placebo arms, respectively. Mean greatest increases from baseline in systolic blood pressure on treatment were 24.4 and 19.6 mmHg in the niraparib and placebo arms, respectively. Mean greatest increases from baseline in diastolic blood pressure on treatment were 15.9 and 13.9 mmHg in the niraparib and placebo arms, respectively.

In the NOVA study, mean pulse rate and blood pressure increased over baseline in the niraparib arm relative to the placebo arm at all on-study assessments. Mean greatest increases from baseline in pulse rate on treatment were 24.1 and 15.8 beats/min in the niraparib and placebo arms, respectively. Mean greatest increases from baseline in systolic blood pressure on treatment were 24.5 and 18.3 mmHg in the niraparib and placebo arms, respectively. Mean greatest increases from baseline in diastolic blood pressure on treatment were 16.5 and 11.6 mmHg in the niraparib and placebo arms, respectively.

#### Cardiac Electrophysiology

The potential for QTc prolongation with niraparib was evaluated in a randomized, placebo-controlled trial in patients with cancer (367 patients on niraparib and 179 patients on placebo). No large changes in the mean QTc interval (>20 ms) were detected in the trial following the treatment of niraparib 300 mg once daily.

### 12.3 Pharmacokinetics

Following a single-dose administration of 300 mg niraparib, the mean ( $\pm$ SD) peak plasma concentration ( $C_{\max}$ ) was 804 ( $\pm$ 403) ng/mL. The exposure ( $C_{\max}$  and AUC) of niraparib increased in a dose-proportional manner with daily doses ranging from 30 mg (0.1 times the approved recommended dose) to 400 mg (1.3 times the approved recommended dose). The accumulation ratio of niraparib exposure following 21 days of repeated daily doses was approximately 2-fold for doses ranging from 30 to 400 mg.

#### Absorption

The absolute bioavailability of niraparib is approximately 73%. Following oral administration of niraparib, peak plasma concentration,  $C_{\max}$ , is reached within 3 hours.

Concomitant administration of a high-fat meal (800 to 1,000 calories with approximately 50% of total caloric content of the meal from fat) did not significantly affect the pharmacokinetics of niraparib.

#### Distribution

Niraparib is 83.0% bound to human plasma proteins. The average ( $\pm$ SD) apparent volume of distribution (Vd/F) was 1,220 ( $\pm$ 1,114) L. In a population pharmacokinetic analysis, the Vd/F of niraparib was 1,074 L in patients with cancer .

#### Elimination

Following multiple daily doses of 300 mg of niraparib, the mean half-life ( $t_{1/2}$ ) is 36 hours. In a population pharmacokinetic analysis, the apparent total clearance (CL/F) of niraparib was 16.2 L/h in patients with cancer.

*Metabolism:* Niraparib is metabolized by carboxylesterases (CEs) to form a major inactive metabolite, which subsequently undergoes glucuronidation.

*Excretion:* Following administration of a single oral 300-mg dose of radio-labeled niraparib, the average percent recovery of the administered dose over 21 days was 47.5% (range: 33.4% to 60.2%) in urine and 38.8% (range: 28.3% to 47.0%) in feces. In pooled samples collected over 6 days, unchanged niraparib accounted for 11% and 19% of the administered dose recovered in urine and feces, respectively.

#### Specific Populations

Age (18 to 65 years), race/ethnicity, and mild to moderate renal impairment ( $CL_{cr} \geq 30$  to 90 mL/min) had no clinically significant effect on the pharmacokinetics of niraparib.

The effect of severe renal impairment ( $CL_{cr} < 30$  mL/min) or end-stage renal disease undergoing hemodialysis on the pharmacokinetics of niraparib is unknown.

*Patients with Hepatic Impairment:* Mild hepatic impairment (total bilirubin <1.5 x ULN and any AST level or bilirubin ≤ ULN and AST > ULN) had no clinically significant effect on the pharmacokinetics of niraparib.

In a trial of patients with moderate hepatic impairment (total bilirubin ≥1.5 x ULN to 3.0 x ULN and any AST level) (n = 8), niraparib AUC<sub>inf</sub> was 1.56 (90% CI: 1.06 to 2.30) times higher compared with patients with normal hepatic function (n = 9) following administration of a single 300-mg dose. Niraparib dosage reduction is recommended for patients with moderate hepatic impairment [see *Dosage and Administration (2.4)*]. Moderate hepatic impairment did not have an effect on niraparib C<sub>max</sub> or on niraparib protein binding.

The effect of severe hepatic impairment (total bilirubin >3.0 x ULN and any AST level) on the pharmacokinetics of niraparib is unknown.

#### Drug Interaction Studies

No clinical drug interaction studies have been performed with ZEJULA.

*In Vitro Studies: Inhibition of Cytochrome P450 (CYP) Enzymes:* Neither niraparib nor the major primary metabolite M1 is an inhibitor of CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, and CYP3A4.

*Induction of CYP Enzymes:* Neither niraparib nor M1 is a CYP3A4 inducer. Niraparib weakly induces CYP1A2 in vitro.

*Substrate of CYP Enzymes:* Niraparib is a substrate of CEs and the resulting M1 is further metabolized through the formation of glucuronides in vivo.

*Inhibition of Uridine 5'-Diphospho-Glucuronosyltransferases (UGTs):* Niraparib did not exhibit inhibitory effect against the UGT isoforms (UGT1A1, UGT1A4, UGT1A9, and UGT2B7) up to 200 microM in vitro. Therefore, the potential for a clinically relevant inhibition of UGTs by niraparib is minimal.

*Inhibition of Transporter Systems:* Niraparib is a weak inhibitor of breast cancer resistance protein (BCRP), but does not inhibit P-glycoprotein (P-gp), bile salt export pump (BSEP), or multidrug resistance-associated protein 2 (MRP2).

Niraparib is an inhibitor of multidrug and toxin extrusion (MATE) 1 and 2 with IC<sub>50</sub> of 0.18 microM and ≤0.14 microM, respectively. Increased plasma concentrations of coadministered drugs that are substrates of these transporters (e.g., metformin) cannot be excluded.

The M1 metabolite is not an inhibitor of P-gp, BCRP, BSEP, MRP2, or MATE1 or 2. Neither niraparib nor M1 is an inhibitor of organic anion transporting polypeptide (OATP)1B1, OATP1B3, organic cation transporter (OCT1)1, organic anion transporter (OAT)1, OAT3, or OCT2.

*Substrate of Transporter Systems:* Niraparib is a substrate of P-gp and BCRP. Niraparib is not a substrate of BSEP, MRP2, or MATE1 or 2. The M1 metabolite is not a substrate of P-gp, BCRP, BSEP, or MRP2. However, M1 is a substrate of MATE1 and 2. Neither niraparib nor M1 is a substrate of OATP1B1, OATP1B3, OCT1, OAT1, OAT3, or OCT2.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Carcinogenicity studies have not been conducted with niraparib.

Niraparib was clastogenic in an in vitro mammalian chromosomal aberration assay and in an in vivo rat bone marrow micronucleus assay. This clastogenicity is consistent with genomic instability resulting from the primary pharmacology of niraparib and indicates potential for genotoxicity in humans. Niraparib was not mutagenic in a bacterial reverse mutation assay (Ames) test.

Fertility studies in animals have not been conducted with niraparib. In repeat-dose oral toxicity studies, niraparib was administered daily for up to 3 months' duration in rats and dogs. Reduced sperm, spermatids, and germ cells in epididymides and testes were observed at doses  $\geq 10$  mg/kg and  $\geq 1.5$  mg/kg in rats and dogs, respectively. These dose levels resulted in systemic exposures approximately 0.3 and 0.012 times, respectively, the human exposure ( $AUC_{0-24h}$ ) at the recommended dose of 300 mg daily. There was a trend toward reversibility of these findings 4 weeks after dosing was stopped.

### **13.2 Animal Toxicology and/or Pharmacology**

In vitro, niraparib bound to DAT, NET, and SERT and inhibited uptake of norepinephrine and dopamine in cells with  $IC_{50}$  values that were lower than the  $C_{min}$  at steady-state in patients receiving the recommended dose. Niraparib has the potential to cause effects in patients related to inhibition of these transporters (e.g., cardiovascular, central nervous system).

Intravenous administration of niraparib to vagotomized dogs over 30 minutes at 1, 3, and 10 mg/kg resulted in an increased range of arterial pressures of 13% to 20%, 18% to 27%, and 19% to 25%, respectively, and increased range of heart rates of 2% to 11%, 4% to 17%, and 12% to 21%, respectively, above pre-dose levels. The unbound plasma concentrations of niraparib in dogs at these dose levels were approximately 0.5, 1.5, and 5.8 times the unbound  $C_{max}$  at steady-state in patients receiving the recommended dose.

In addition, niraparib crossed the blood-brain barrier in rats and monkeys following oral administration. The cerebrospinal fluid:plasma  $C_{max}$  ratios of niraparib administered at 10 mg/kg orally to 2 rhesus monkeys were 0.10 and 0.52.

## 14 CLINICAL STUDIES

### 14.1 First-Line Maintenance Treatment of Advanced Ovarian Cancer

PRIMA (NCT02655016) was a double-blind, placebo-controlled trial in which patients (N = 733) in complete or partial response to first-line platinum-based chemotherapy were randomized 2:1 to ZEJULA or matched placebo. Initially, the patients received a starting dosage of 300 mg once daily regardless of body weight or platelet count. The study was amended to include a starting dose of 200 mg for patients weighing <77 kg (<170 lbs) OR with a platelet count of <150,000/mcL or 300 mg for patients weighing ≥77 kg (≥170 lbs) AND who had a platelet count ≥150,000/mcL.

Patients were randomized post-completion of first-line platinum-based chemotherapy plus surgery. Randomization was stratified by best response during the front-line platinum regimen (complete response vs. partial response), neoadjuvant chemotherapy (NACT) (yes vs. no), and HRD status (positive vs. negative or not determined). HRD status was determined using the FDA-approved Myriad myChoice CDx assay. HRD positive status included either tumor *BRCA* mutant (*tBRCAm*) or a genomic instability score (GIS) ≥42.

The major efficacy outcome measure, progression-free survival (PFS), was determined by blinded independent central review (BICR) per Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1. In some cases, criteria other than RECIST, such as clinical signs and symptoms and increasing CA-125, were also applied. Overall survival was an additional efficacy outcome measure. PFS testing was performed hierarchically: first in the homologous recombination (HR)-deficient (HRD positive) population, then in the overall population. The median age of 62 ranged from 32 to 85 years among patients randomized with ZEJULA and 33 to 88 years among patients randomized with placebo. Eighty-nine percent of all patients were White. Sixty-nine percent of patients randomized with ZEJULA and 71% of patients randomized with placebo had an Eastern Cooperative Oncology Group Performance Status (ECOG PS) of 0 at study baseline. Approximately 45% of patients were enrolled in the U.S. or Canada. In the overall population, 65% of patients had stage III disease and 35% had stage IV disease. Sixty-seven percent of the patients received NACT. Sixty-nine percent of the patients had a complete response to the first-line platinum-based chemotherapy. Approximately 35% (n = 258) of patients received a starting dose of 200 or 300 mg depending on baseline body weight and platelet count. Among those patients, 186 patients received a starting dose of 200 mg.

PRIMA demonstrated a statistically significant improvement in PFS for patients randomized to ZEJULA as compared with placebo in the HR-deficient and overall population (Table 12, Figure 1, and Figure 2).



**Table 12. Efficacy Results–PRIMA (determined by BICR<sup>a</sup>)**

	HR-Deficient Population		Overall Population	
	ZEJULA (n = 247)	Placebo (n = 126)	ZEJULA (n = 487)	Placebo (n = 246)
Progression-free survival events, n (%)	81 (33)	73 (58)	232 (48)	155 (63)
Progression-free survival median in months (95% CI)	21.9 (19.3, NE)	10.4 (8.1, 12.1)	13.8 (11.5, 14.9)	8.2 (7.3, 8.5)
Hazard ratio <sup>b</sup> (95% CI)	0.43 (0.31, 0.59)		0.62 (0.50, 0.76)	
<i>P</i> value <sup>c</sup>	<0.0001		<0.0001	

HR = Homologous Recombination, NE = not estimable.


<sup>a</sup> Efficacy analysis was based on blinded independent central review.

<sup>b</sup> Based on a stratified Cox proportional hazards model.


<sup>c</sup> Based on a stratified log-rank test.

In exploratory subgroup analyses of patients who were administered a starting dose of ZEJULA or matched placebo based on baseline weight or platelet count, the hazard ratio for PFS was 0.39 (95% CI [0.22, 0.72]) in the HR-deficient subgroup (n = 130) and 0.68 (95% CI [0.48, 0.97]) in the overall population (n = 258).

**Figure 1. Progression-Free Survival in Patients with HR-Deficient Tumors (Intent-to-Treat Population, n = 373)**



**Figure 2. Progression-Free Survival in the Overall Population (Intent-to-Treat Population, n = 733)**



At the time of the PFS analysis, overall survival data were immature, with 11% deaths in the overall population.

## 14.2 Maintenance Treatment of Recurrent Ovarian Cancer

NOVA (NCT01847274) was a double-blind, placebo-controlled trial in which patients (N = 553) with platinum-sensitive recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer were randomized 2:1 to ZEJULA 300 mg orally daily or matched placebo within 8 weeks of the last therapy. Treatment was continued until disease progression or unacceptable toxicity. All patients had received at least 2 prior platinum-containing regimens and were in response (complete or partial) to their most recent platinum-based regimen.

Randomization was stratified by time to progression after the penultimate platinum therapy (6 to <12 months and  $\geq 12$  months), use of bevacizumab in conjunction with the penultimate or last platinum regimen (yes/no), and best response during the most recent platinum regimen (complete response and partial response). Eligible patients were assigned to 1 of 2 cohorts based on the results of the BRACAnalysis CDx. Patients with deleterious or suspected deleterious germline *BRCA* mutations (g*BRCA*m) were assigned to the germline *BRCA*-mutated (g*BRCA*mut) cohort (n = 203), and those without germline *BRCA* mutations were assigned to the non-g*BRCA*mut cohort (n = 350).

The major efficacy outcome measure, PFS, was determined primarily by central independent assessment per RECIST version 1.1. In some cases, criteria other than RECIST, such as clinical signs and symptoms and increasing CA-125, were also applied.

The median age of patients ranged from 57 to 64 years among patients treated with ZEJULA and 58 to 67 years among patients treated with placebo. Eighty-six percent of all patients were White. Sixty-seven percent of patients receiving ZEJULA and 69% of patients receiving placebo had an ECOG PS of 0 at study baseline. Approximately 40% of patients were enrolled in the U.S. or Canada, and 51% of all patients were in complete response to most recent platinum-based regimen, with 39% on both arms with an interval of 6 to 12 months since the penultimate platinum regimen. Twenty-six percent of those treated with ZEJULA and 31% treated with placebo had received prior bevacizumab therapy. Approximately 40% of patients had 3 or more lines of treatment.

The trial demonstrated a statistically significant improvement in PFS for patients randomized to ZEJULA as compared with placebo in the *gBRCAmut* cohort and the non-*gBRCAmut* cohort (Table 13, Figure 3, and Figure 4).

**Table 13. Efficacy Results–NOVA (IRC Assessment,<sup>a</sup> Intent-to-Treat Population)**

	<i>gBRCAmut</i> Cohort		non- <i>gBRCAmut</i> Cohort	
	<b>ZEJULA (n = 138)</b>	<b>Placebo (n = 65)</b>	<b>ZEJULA (n = 234)</b>	<b>Placebo (n = 116)</b>
Progression-free survival median in months (95% CI)	21.0 (12.9, NR)	5.5 (3.8, 7.2)	9.3 (7.2, 11.2)	3.9 (3.7, 5.5)
Hazard ratio <sup>b</sup> (95% CI)	0.26 (0.17, 0.41)		0.45 (0.34, 0.61)	
<i>P</i> value <sup>c</sup>	<0.0001		<0.0001	


IRC = Independent Review Committee, *gBRCAmut* = germline *BRCA*-mutated, NR = not reached.

<sup>a</sup> Efficacy analysis was based on blinded central independent radiologic and clinical oncology review committee.

<sup>b</sup> Based on a stratified Cox proportional hazards model.


<sup>c</sup> Based on a stratified log-rank test.

**Figure 3. Progression-Free Survival in the *gBRCA*mut Cohort Based on IRC Assessment (Intent-to-Treat Population, n = 203)**



*gBRCA*mut = germline *BRCA*-mutated, IRC = Independent Review Committee.

**Figure 4. Progression-Free Survival in the Non-*gBRCA*mut Cohort Overall Based on IRC Assessment (Intent-to-Treat Population, n = 350)**



*gBRCA*mut = germline *BRCA*-mutated, IRC = Independent Review Committee.

At the time of the PFS analysis, limited overall survival data were available with 17% deaths across the 2 cohorts.

### 14.3 Treatment of Advanced Ovarian Cancer after 3 or More Chemotherapies

The efficacy of ZEPJULA was studied in 98 patients with advanced ovarian cancer with HRD positive tumors in the single-arm QUADRA (NCT02354586) trial. Patients were required to have been treated with 3 or more prior lines of chemotherapy and those with prior exposure to PARP inhibitors were excluded. Patients were selected using a clinical trial assay. Those without *BRCA* mutations must have progressed at least 6 months after their last dose of platinum therapy. All patients received ZEPJULA capsules at a starting dosage of 300 mg once daily as monotherapy until disease progression or unacceptable toxicity.

HRD positive status was determined using the Myriad myChoice CDx as either *tBRCAm* (n = 63) and/or a GIS  $\geq 42$  (n = 35). GIS is an algorithmic measurement of loss of heterozygosity, telomeric allelic imbalance, and large-scale state transitions.

The major efficacy outcome measures were objective response rate (ORR) and duration of response as assessed by the investigator according to RECIST v. 1.1.

The median age of the patients was 63 years (range: 39 to 91 years), the majority were White (82%), and all had an ECOG PS of 0 (59%) or 1 (41%).

The efficacy results for QUADRA are summarized in Table 14.

**Table 14. Efficacy Results–QUADRA (Investigator Assessment)**

Efficacy Results	HRD Positive Cohort <sup>a</sup> (N = 98)
Objective response rate (95% CI) <sup>b</sup>	24% (16, 34)
Complete responses	0%
Partial responses	24%
Median duration of response in months (95% CI)	8.3 (6.5, NE)

NE = not estimable.

<sup>a</sup> Homologous recombination deficiency (HRD) positive status is defined as *tBRCA*-mutated and/or genomic instability score  $\geq 42$ .

<sup>b</sup> Confirmed response rate. The objective response rate as assessed by blinded independent central review was consistent.

For patients with *tBRCAm* ovarian cancer, investigator-assessed ORR was 39% (7/18; 95% CI: [17, 64]) in patients with platinum-sensitive disease, 29% (6/21; 95% CI: [11, 52]) in patients with platinum-resistant disease, and 19% (3/16; 95% CI: [4, 46]) in patients with platinum-refractory disease.

For patients with platinum-sensitive GIS-positive disease (without *BRCA*mut) (n = 35), investigator-assessed ORR was 20% (95% CI [8, 37]).

## 16 HOW SUPPLIED/STORAGE AND HANDLING

ZEJULA is available as capsules having a white body printed with “100 mg” in black ink, and a purple cap printed with “Niraparib” in white ink.

Each capsule contains 100 mg of niraparib free base.

ZEJULA capsules are packaged as

90-count bottles                      NDC 69656-103-90

30-count bottles                      NDC 69656-103-30

Store at 20°C to 25°C (68°F to 77°F); excursions are permitted between 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

## **17 PATIENT COUNSELING INFORMATION**

Advise the patient to read the FDA-approved patient labeling (Patient Information).

### Myelodysplastic Syndrome/Acute Myeloid Leukemia

Advise patients to contact their healthcare provider if they experience weakness, feeling tired, fever, weight loss, frequent infections, bruising, bleeding easily, breathlessness, blood in urine or stool, and/or laboratory findings of low blood cell counts or a need for blood transfusions. This may be a sign of hematological toxicity or MDS or AML, which has been reported in patients treated with ZEJULA [see *Warnings and Precautions (5.1)*].

### Bone Marrow Suppression

Advise patients that periodic monitoring of their blood counts is required. Advise patients to contact their healthcare provider for new onset of bleeding, fever, or symptoms of infection [see *Warnings and Precautions (5.2)*].

### Hypertension and Cardiovascular Effects

Advise patients to undergo blood pressure and heart rate monitoring at least weekly for the first 2 months, then monthly for the first year of treatment and periodically thereafter. Advise patients to contact their healthcare provider if blood pressure is elevated [see *Warnings and Precautions (5.3)*].

### Posterior Reversible Encephalopathy Syndrome

Inform patients that they are at risk of developing posterior reversible encephalopathy syndrome (PRES) that can present with signs and symptoms including seizure, headaches, altered mental status, or vision changes. Advise patients to contact their healthcare provider if they develop any of these signs or symptoms [see *Warnings and Precautions (5.4)*].

### Dosing Instructions

Inform patients on how to take ZEJULA [see *Dosage and Administration (2.2)*]. ZEJULA should be taken once daily. Instruct patients that if they miss a dose of ZEJULA not to take an extra dose to make up for the one that they missed. They should take their next dose at the regularly scheduled time. Each capsule should be swallowed whole. ZEJULA may be taken with or without food. Bedtime administration may be a potential method for managing nausea.

### Embryo-Fetal Toxicity

Advise females to inform their healthcare provider if they are pregnant or become pregnant. Inform female patients of the risk to a fetus and potential loss of the pregnancy [see *Warnings and Precautions (5.5), Use in Specific Populations (8.1)*].



### Contraception

Advise females of reproductive potential to use effective contraception during treatment with ZEPJULA and for at least 6 months after receiving the last dose [see *Use in Specific Populations (8.3)*].

### Lactation

Advise patients not to breastfeed while taking ZEPJULA and for 1 month after the last dose [see *Use in Specific Populations (8.2)*].

### Allergic Reactions to FD&C Yellow No. 5 (Tartrazine)

Advise patients that ZEPJULA capsules contain FD&C Yellow No. 5 (tartrazine), which may cause allergic-type reactions (including bronchial asthma) in certain susceptible persons or in patients who also have aspirin hypersensitivity [see *Warnings and Precautions (5.6)*].

Trademarks are owned by or licensed to the GSK group of companies.

Manufactured for



GlaxoSmithKline  
Research Triangle Park, NC 27709

©2021 GSK group of companies or its licensor.

ZJC:1PI

**PHARMACIST—DETACH HERE AND GIVE LEAFLET TO PATIENT**

---

**Patient Information**  
**ZEJULA (zuh-JOO-luh)**  
**(niraparib)**  
**capsules**

**What is the most important information I should know about ZEJULA?**

**ZEJULA may cause serious side effects including:**

- **Bone marrow problems called myelodysplastic syndrome (MDS) or a type of cancer of the blood called acute myeloid leukemia (AML).** Some people who have ovarian cancer and who have received previous treatment with chemotherapy or certain other medicines for their cancer have developed MDS or AML during treatment with ZEJULA. MDS or AML may lead to death. If you develop MDS or AML, your healthcare provider will stop treatment with ZEJULA.

Symptoms of low blood cell counts (low red blood cells, low white blood cells, and low platelets) are common during treatment with ZEJULA, but can be a sign of serious bone marrow problems, including MDS or AML. Symptoms may include:

- weakness
- feeling tired
- weight loss
- frequent infections
- fever
- shortness of breath
- blood in urine or stool
- bruising or bleeding more easily

Your healthcare provider will do blood tests to check your blood cell counts:

- before treatment with ZEJULA.
  - weekly for the first month of treatment with ZEJULA.
  - every month for the next 11 months, then as needed during treatment with ZEJULA.
- **High blood pressure.** High blood pressure is common during treatment with ZEJULA and can become serious. Your healthcare provider will check your blood pressure and heart rate at least weekly for the first 2 months, then monthly for the first year and as needed thereafter during your treatment with ZEJULA.
  - **Posterior reversible encephalopathy syndrome (PRES).** PRES is a condition that affects the brain and may happen during treatment with ZEJULA. If you have headache, vision changes, confusion, or seizure with or without high blood pressure, please contact your healthcare provider.

**See “What are the possible side effects of ZEJULA?” for more information about side effects.**

**What is ZEJULA?**

ZEJULA is a prescription medicine used for the:

- maintenance treatment of adults with advanced ovarian cancer, fallopian tube cancer, or primary peritoneal cancer. ZEJULA is used after the cancer has responded (complete or partial response) to treatment with platinum-based chemotherapy.
- maintenance treatment of adults with ovarian cancer, fallopian tube cancer, or primary peritoneal cancer that comes back. ZEJULA is used after the cancer has responded (complete or partial response) to treatment with platinum-based chemotherapy.

- treatment of adults with advanced ovarian cancer, fallopian tube cancer, or primary peritoneal cancer who have been treated with 3 or more prior types of chemotherapy and who have tumors with:
  - a certain *BRCA* gene mutation, **or**
  - gene mutation problems and who have progressed more than 6 months after their last treatment with platinum-based chemotherapy.

Your healthcare provider will perform a test to make sure that ZEPJULA is right for you.

It is not known if ZEPJULA is safe and effective in children.

**Before taking ZEPJULA, tell your healthcare provider about all of your medical conditions, including if you:**

- have heart problems.
- have liver problems.
- have high blood pressure.
- are allergic to FD&C Yellow No. 5 (tartrazine) or aspirin. ZEPJULA capsules contain FD&C Yellow No. 5 (tartrazine), which may cause allergic-type reactions (including bronchial asthma) in certain people, especially people who also have an allergy to aspirin.
- are pregnant or plan to become pregnant. ZEPJULA can harm your unborn baby and may cause loss of pregnancy (miscarriage).
  - If you are able to become pregnant, your healthcare provider may perform a pregnancy test before you start treatment with ZEPJULA.
  - Females who are able to become pregnant should use effective birth control (contraception) during treatment with ZEPJULA and for 6 months after the last dose of ZEPJULA. Talk to your healthcare provider about birth control methods that may be right for you.
  - Tell your healthcare provider right away if you become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if ZEPJULA passes into your breast milk. Do not breastfeed during treatment with ZEPJULA and for 1 month after the last dose of ZEPJULA. Talk to your healthcare provider about the best way to feed your baby during this time.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**How should I take ZEPJULA?**

- Take ZEPJULA exactly as your healthcare provider tells you to.
- Take ZEPJULA 1 time each day, at the same time each day.
- ZEPJULA may be taken with or without food.
- ZEPJULA capsules should be swallowed whole. Do not chew, crush, or split ZEPJULA capsules before swallowing.
- Taking ZEPJULA at bedtime may help relieve any nausea symptoms you may have.
- Do not stop taking ZEPJULA without first talking with your healthcare provider.
- If you miss a dose of ZEPJULA, take your next dose at your scheduled time. Do not take an extra dose to make up for a missed dose.

- If you vomit after taking a dose of ZEPJULA, do not take an extra dose. Take your next dose at your scheduled time.
- If you take too much ZEPJULA, call your healthcare provider or go to the nearest hospital emergency room right away.

**What are the possible side effects of ZEPJULA?**

**ZEPJULA may cause serious side effects, including:**

- See “What is the most important information I should know about ZEPJULA?”

The most common side effects of ZEPJULA include:

- |                               |  |
|-------------------------------|--|
| • heart not beating regularly | • changes in liver function or other blood tests |
| • nausea                      | • pain in your muscles and back                  |
| • constipation                | • headache                                       |
| • vomiting                    | • dizziness                                      |
| • pain in the stomach area    | • change in the way food tastes                  |
| • mouth sores                 | • trouble sleeping                               |
| • diarrhea                    | • anxiety  |
| • indigestion or heartburn    | • sore throat                                    |
| • dry mouth                   | • shortness of breath                            |
| • tiredness                   | • cough  |
| • loss of appetite            | • rash   |
| • urinary tract infection     | • changes in the amount or color of your urine   |

Your healthcare provider may change your dose, temporarily stop treatment, or permanently stop treatment with ZEPJULA if you have certain side effects.

These are not all the possible side effects of ZEPJULA.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store ZEPJULA?**

Store ZEPJULA at room temperature between 68° to 77°F (20° to 25°C).

**Keep ZEPJULA and all medicines out of the reach of children.**

**General information about the safe and effective use of ZEPJULA.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ZEPJULA for a condition for which it was not prescribed. Do not give ZEPJULA to other people, even if they have the same symptoms that you have. It may harm them. You can ask your healthcare provider or pharmacist for information about ZEPJULA that is written for health professionals.

**What are the ingredients in ZEPJULA?**

**Active ingredient:** niraparib.

**Inactive ingredients:**

Capsule fill: magnesium stearate and lactose monohydrate.

Capsule shell: titanium dioxide and gelatin in the white capsule body and FD&C Blue No. 1, FD&C Red No. 3, FD&C Yellow No. 5 (tartrazine), and gelatin in the purple capsule cap.

The black printing ink: shellac, dehydrated alcohol, isopropyl alcohol, butyl alcohol, propylene glycol, purified water, strong ammonia solution, potassium hydroxide, and black iron oxide.

The white printing ink: shellac, dehydrated alcohol, isopropyl alcohol, butyl alcohol, propylene glycol, sodium hydroxide, povidone, and titanium dioxide.



For more information about ZEJULA, call 1-888-825-5249.

Trademarks are owned by or licensed to the GSK group of companies.

Manufactured for:

GlaxoSmithKline, Research Triangle Park, NC 27709

©2021 GSK group of companies or its licensor.

ZJC:1PIL

This Patient Information has been approved by the U.S. Food and Drug Administration

Revised: March 2021