

## Efectos adversos\*

Acontecimiento	Niraparib (n =367)		Placebo (n =179)	
	Cualquier grado	Grado 3 o4	Cualquier grado	Grado 3 o 4
	Número de pacientes (porcentaje)			
Náuseas	270 (73,6)	11 (3,0)	63 (35,2)	2 (1,1)
Trombocitopenia †	225 (61,3)	124 (33,8)	10 (5,6)	1 (0,6)
Fatiga‡	218 (59,4)	30 (8,2)	74 (41,3)	1 (0,6)
Anemia§	184 (50,1)	93 (25,3)	12 (6,7)	0
Estreñimiento	146 (39,8)	2 (0,5)	36 (20,1)	1 (0,6)
Vómitos	126 (34,3)	7 (1,9)	29 (16,2)	1 (0,6)
Neutropenia¶	111 (30,2)	72 (19,6)	11 (6,1)	3 (1,7)
Cefalea	95 (25,9)	1 (0,3)	17 (9,5)	0
Disminución del apetito	93 (25,3)	1 (0,3)	26 (14,5)	1 (0,6)
Insomnio	89 (24,3)	1 (0,3)	13 (7,3)	0
Dolor abdominal	83 (22,6)	4 (1,1)	53 (29,6)	3 (1,7)
Disnea	71 (19,3)	4 (1,1)	15 (8,4)	2 (1,1)
Hipertensión	71 (19,3)	30 (8,2)	8 (4,5)	4 (2,2)
Diarrea	70 (19,1)	1 (0,3)	37 (20,7)	2 (1,1)
Mareo	61 (16,6)	0	13 (7,3)	0
Tos	55 (15,0)	0	8 (4,5)	0
Dolor de espalda	49 (13,4)	2 (0,5)	21 (11,7)	0
Artralgia	43 (11,7)	1 (0,3)	22 (12,3)	0
Dispepsia	42 (11,4)	0	17 (9,5)	0
Nasofaringitis	41 (11,2)	0	13 (7,3)	0
Infección del tracto urinario	38 (10,4)	3 (0,8)	11 (6,1)	2 (1,1)
Palpitaciones	38 (10,4)	0	3 (1,7)	0
Disgeusia	37 (10,1)	0	7 (3,9)	0
Mialgia	30 (8,2)	1 (0,3)	18 (10,1)	0
Distensión abdominal	28 (7,6)	0	22 (12,3)	1 (0,6)